

DAFTAR PUSTAKA

1. Augustin R, Carayannopoulos MO, Dowd LO, Phay JE, Moley JF, Moley KH. Identification and characterization of human glucose transporter-like protein-9 (GLUT9): alternative splicing alters trafficking. *Journal of Biological Chemistry*. 2004;279(16):16229-16236.
2. Manolescu AR, Augustin R, Moley K, Cheeseman C. A highly conserved hydrophobic motif in the exofacial vestibule of fructose transporting SLC2A proteins acts as a critical determinant of their substrate selectivity. *Molecular membrane biology*. 2007;24(5-6):455-463.
3. Manolescu AR, Witkowska K, Kinnaird A, Cessford T, Cheeseman C. Facilitated hexose transporters: new perspectives on form and function. *Physiology*. 2007;22(4):234-240.
4. Caulfield MJ, Munroe PB, O'Neill D, Witkowska K, Charchar FJ, Doblado M, Evans S, Eyheramendy S, Onipinla A, Howard P, Shaw-Hawkins S. SLC2A9 is a high-capacity urate transporter in humans. *PLoS medicine*. 2008;5(10):e197.
5. Anzai N, Ichida K, Jutabha P, Kimura T, Babu E, Jin CJ, Srivastava S, Kitamura K, Hisatome I, Endou H, Sakurai H. Plasma urate level is directly regulated by a voltage-driven urate efflux transporter URATv1 (SLC2A9) in humans. *Journal of Biological Chemistry*. 2008;283(40):26834-26838. *Journal of Biological Chemistry*. 2008;283(46):32152-32152.
6. Kimura T, Amonpatumrat S, Tsukada A, Fukutomi T, Jutabha P, Thammapatip T, Lee EJ, Ichida K, Anzai N, Sakurai H. Increased expression of SLC2A9 decreases urate excretion from the kidney. *Nucleosides, Nucleotides and Nucleic Acids*. 2011;30(12):1295-1301.
7. Johnson RJ, Tittle S, Cade JR, et al. Uric acid, evolution and primitive cultures. *Semin Nephrol*. 2005;25:3-8.

8. Conen D, Wietlisbach V, Bovet P, et al. Prevalence of hyperuricemia and relation of serum uric acid with cardiovascular risk factors in a developing country. *BMC Public Health*. 2004;4:9.
9. Klein BE, Klein R, Lee KE. Components of the metabolic syndrome and risk of cardiovascular disease and diabetes in beaver dam. *Diabetes Care*. 2002;25:1790–1794.
10. Bos MJ, Koudstaal PJ, Hofman A, et al. Uric acid is a risk factor for myocardial infarction and stroke: the Rotterdam study. *Stroke*. 2006;37:1503–1507.
11. Dehghan A, van Hoek M, Sijbrands EJ, et al. High serum uric acid as a novel risk factor for type 2 diabetes. *Diabetes Care*. 2008;31:361–362.
12. Rosolowsky ET, Ficociello LH, Maselli NJ, et al. High-normal serum uric acid is associated with impaired glomerular filtration rate in nonproteinuric patients with type 1 diabetes. *Clin J Am Soc Nephrol*. 2008;3:706–713.
13. Weiner DE, Tighiouart H, Elsayed EF, et al. Uric acid and incident kidney disease in the community. *J Am Soc Nephrol*. 2008;19(6):1204-1211.
14. Chonchol M, Shlipak MG, Katz R, et al. Relationship of uric acid with progression of kidney disease. *Am J Kidney Dis*. 2007;50:239–247.
15. Caulfield MJ, Munroe PB, O'Neill D, Witkowska K, Charchar FJ, Doblado M, Evans S, Eyheramendy S, Onipinla A, Howard P, Shaw-Hawkins S. SLC2A9 is a high-capacity urate transporter in humans. *PLoS medicine*. 2008;5(10):e197.
16. Li, S., Sanna, S., Maschio, A. et al. The GLUT9 gene is associated with serum uric acid levels in Sardinia and Chianti cohorts. *PLoS Genet*. 2007;3:e194.
17. Wallace, C., Newhouse, S. J., Braund, P. et al. Genomewide association study identifies genes for biomarkers of cardiovascular disease: Serum urate and dyslipidemia. *Am J Hum Genet*. 2008;82:139–149.
18. Matsuo, H., Chiba, T., Nagamori, S. et al. Mutations in glucose transporter 9 gene SLC2A9 cause renal hypouricemia. *Am J Hum Genet*. 2008;83:744–51.

19. Vitart, V., Rudan, I., Hayward, C. et al. SLC2A9 is a newly identified urate transporter influencing serum urate concentration, urate excretion and gout. *Nat Genet.* 2008;40:437–42.
20. Doring, A., Gieger, C., Mehta, D. et al. SLC2A9 influences uric acid concentrations with pronounced sex-specific effects. *Nat Genet.* 2008;40:430–6.
21. Sull, J. W., Park, E. J., Lee, M., & Jee SH. Effects of SLC2A9 variants on uric acid levels in a Korean population. *Rheumatol Int.* 2012;33(1):19–23.
22. Hollis-Moffatt, J.E., Xu, X., Dalbeth, N., Merriman, M.E., Topless, R., Waddell, C., Gow, P.J., Harrison, A.A., Highton, J., Jones, P.B. and Stamp, L.K., Role of the urate transporter SLC2A9 gene in susceptibility to gout in New Zealand Māori, Pacific Island, and Caucasian case–control sample sets. *Arthritis & Rheumatism: Official Journal of the American College of Rheumatology.* 2009; 60(11), pp.3485-3492.
23. Putrohari R.D, Simanjuntak T, Sudoyo H, dkk. Diaspora Melanesia di Nusantara. Direktorat Sejarah dan Nilai Budaya, Direktorat Jenderal Kebudayaan, Kementerian Pendidikan dan Kebudayaan. 2015. Jakarta.
24. Ingman M, Gyllensten U. Mitochondrial genome variation and evolutionary history of Australian and New Guinean aborigines. *Genome research.* 2003;13(7):1600-1606.
25. Van Holst Pellekaan SM, Frommer M, Sved JA, Boettcher B. Mitochondrial control-region sequence variation in aboriginal Australians. *The American Journal of Human Genetics.* 1998;62(2):435-449.
26. Rasmussen M, Guo X, Wang Y, Lohmueller KE, Rasmussen S, Albrechtsen A, Skotte L, Lindgreen S, Metspalu M, Jombart T, Kivisild T. An Aboriginal Australian genome reveals separate human dispersals into Asia. *Science.* 2011;334(6052):94-8.
27. Mona S, Tommaseo-Ponzetta M, Brauer S, Sudoyo H, Marzuki S, Kayser M. Patterns of Y-chromosome diversity intersect with the Trans-New Guinea hypothesis. *Mol Biol Evol.* 2007;24:2546–2555.

28. Friedlaender JS, Friedlaender FR, Reed FA, Kidd KK, Kidd JR, Chambers GK, Lea RA, Loo JH, Koki G, Hodgson JA, Merriwether DA. The genetic structure of Pacific Islanders. *PLoS genetics*. 2008;4(1):e19.
29. Kayser M, Brauer S, Weiss G, Schiefenhover W, Underhill P, Shen P, Oefner P, Tommaseo-Ponzetta M, Stoneking M. Reduced Y-chromosome, but not mitochondrial DNA, diversity in human populations from West New Guinea. *Am J Hum Genet*. 2003;72: 281–302.
30. Bergström A, Oppenheimer SJ, Mentzer AJ, Auckland K, Robson K, Attenborough R, Alpers MP, Koki G, Pomat W, Siba P, Xue Y. A Neolithic expansion, but strong genetic structure, in the independent history of New Guinea. *Science*. 2017 S;357(6356):1160- 1163.
31. Stoneking M, Jorde LB, Bhatia K, Wilson AC. Geographic variation in human mitochondrial DNA from Papua New Guinea. *Genetics*. 1990;124(3):717-733.
32. Maekawa K, Itoda M, Sai K, Saito Y, Kaniwa N, Shirao K, Hamaguchi T, Kunitoh H, Yamamoto N, Tamura T, Minami H. Genetic variation and haplotype structure of the ABC transporter gene ABCG2 in a Japanese population. *Drug metabolism and pharmacokinetics*. 2006;21(2):109-21.
33. Tumonggor MK, Karafet TM, Hallmark B, Lansing JS, Sudoyo H, Hammer MF, Cox MP. The Indonesian archipelago: an ancient genetic highway linking Asia and the Pacific. *Journal of human genetics*. 2013;58(3):165-73.
34. Pagani L, Lawson DJ, Jagoda E, Mörseburg A, Eriksson A, Mitt M, Clemente F, Hudjashov G, DeGiorgio M, Saag L, Wall JD. Genomic analyses inform on migration events during the peopling of Eurasia. *Nature*. 2016;538(7624):238-242.
35. Upton S. The impact of migration on the people of Papua, Indonesia. *Department of History and Philosophy University of New South Wales*. 2009; pp. 70-80.
36. Pritchard JK, Przeworski M. Linkage disequilibrium in humans: models and data. *The American Journal of Human Genetics*. 2001;69(1):1-14.

37. Dalbeth N, Allan J, Gamble GD, Phipps-Green A, Flynn TJ, Mihov B, et al. Influence of genetic variants on renal uric acid handling in response to frusemide: An acute intervention study. *RMD Open*. 2017;3(1):1–8.
38. Wrigley R, Phipps-Green AJ, Topless RK, Major TJ, Cadzow M, Riches P, et al. Pleiotropic effect of the ABCG2 gene in gout: Involvement in serum urate levels and progression from hyperuricemia to gout. *Arthritis Res Ther*. 2020;22(1):1–10.
39. Batt C, Phipps-Green AJ, Black MA, Cadzow M, Merriman ME, Topless R, Gow P, Harrison A, Highton J, Jones P, Stamp L. Sugar-sweetened beverage consumption: a risk factor for prevalent gout with SLC2A9 genotype-specific effects on serum urate and risk of gout. *Annals of the rheumatic diseases*. 2014;73(12):2101-2106.
40. Dalbeth N, House ME, Gamble GD, Horne A, Pool B, Purvis L, Stewart A, Merriman M, Cadzow M, Phipps-Green A, Merriman TR. Population-specific influence of SLC2A9 genotype on the acute hyperuricaemic response to a fructose load. *Annals of the rheumatic diseases*. 2013;72(11):1868-73.
41. World-first genomic study of Aboriginal Australians. 22 September 2016. <https://news.griffith.edu.au/2016/09/22world-first-genomic-study-of-aboriginal-australians/>
42. Xu S, Pugach I, Stoneking M, Kayser M, Jin L, HUGO Pan-Asian SNP Consortium. Genetic dating indicates that the Asian–Papuan admixture through Eastern Indonesia corresponds to the Austronesian expansion. *Proceedings of the National Academy of Sciences*. 2012;109(12):4574-4579.
43. Karafet TM, Hallmark B, Cox MP, Sudoyo H, Downey S, Lansing JS, Hammer MF. Major east–west division underlies Y chromosome stratification across Indonesia. *Molecular biology and evolution*. 2010;27(8):1833-44.
44. Kayser M, Brauer S, Cordaux R, et al. (15 co-authors). Melanesian and Asian origins of Polynesians: mtDNA and Y chromosome gradients across the Pacific. *Mol Biol Evol*. 2006;23:2234–2244.

45. Tefu, M.O. and Karwur, F.F. Pitarah Manusia Nusa Tenggara Timur Berdasarkan Ceritera Kromosom Y. *Scientiae Educatia: Jurnal Pendidikan Sains*. 2017;6(2), pp.144-165.

