

DAFTAR PUSTAKA

- [1] S. Budiman, H. H. Bahti, A. Mutalib, and A. Anggraen, "Pemisahan Gadolinium Sebagai Contrast Agent Pada Mri (Magnetic Resonance Imaging) Dengan Ligan Asam Di-(2-Etilheksil) Fosfat," *J. Sains dan Kesehat.*, vol. 1, no. 9, pp. 510–516, 2017, doi: <https://doi.org/10.25026/jsk.v1i9.88>.
- [2] S. T. Bird *et al.*, "First-Trimester Exposure to Gadolinium-based Contrast Agents: A Utilization Study of 4.6 Million U.S. Pregnancies," *Radiology*, vol. 293, no. 1, pp. 193–200, Oct. 2019, doi: <https://doi.org/10.1148/radiol.2019190563>.
- [3] A. Silvio and C. Peter, "Biodistribution of gadolinium-based contrast agents, including gadolinium deposition," *J. Magn. Reson. Imaging*, vol. 30, no. 6, pp. 1259–1267, 2009, doi: DOI: 10.1002/jmri.21969.
- [4] L. Chehabeddine, T. Al Saleh, M. Baalbaki, E. Saleh, S. J. Khoury, and S. Hannoun, "Cumulative administrations of gadolinium-based contrast agents: risks of accumulation and toxicity of linear vs macrocyclic agents," *Crit. Rev. Toxicol.*, vol. 49, no. 3, pp. 262–279, 2019, doi: DOI: 10.1080/10408444.2019.1592109.
- [5] L. B. M and S. Wangko, "Peran Sel Kupffer Pada Steatohepatitis Alkohol," *J. Biomedik*, vol. 4, no. 2, 2013, [Online]. Available: <https://ejournal.unsrat.ac.id/index.php/biomedik/article/download/755/12190>
- [6] G. Tapia, F. Troncoso, M. Galleano, V. Fernandez, S. Puntarulo, and L. A. Videla, "Time course study of the influence of acute iron overload on kupffer cell functioning and hepatotoxicity assessed in the isolated perfused rat liver," *Hepatology*, vol. 27, no. 5, pp. 1311–1316, 1998, doi: DOI: 10.1002/hep.510270517.
- [7] E. R. Sari, G. Maslebu, and A. Sutresno, "Studi Difusi Ca²⁺ pada Sinapsis menggunakan Metode Monte Carlo Cell," 2020, [Online]. Available: <https://iptek.its.ac.id/index.php/jfa/article/view/16%281%2910>
- [8] B. Haryanto, "Pengaruh Pemilihan Kondisi Batas, Langkah Ruang, Langkah Waktu, dan Koefisien Difusi pada Model Difusi," *Aplika: Jurnal Ilmu Pengetahuan dan Teknologi*, vol. 8, no. 1, pp. 1–7, 2014.
- [9] L. Telgmann, C. A. Wehe, J. Künnemeyer, A. Bülter, M. Sperling, and U. Karst, "Speciation of Gd-based MRI contrast agents and potential products of transmetalation with iron ions or parenteral iron supplements," pp. 2133–2141, 2012, doi: DOI: 10.1007/s00216-012-6404-x.
- [10] F. Martino, G. Amici, M. Rosner, C. Ronco, and G. Novara, "Gadolinium-based contrast media nephrotoxicity in kidney impairment: The physiological conditions for the perfect murder," *J. Clin. Med.*, vol. 10, no. 2, pp. 1–15, 2021, doi: doi: 10.3390/jcm10020271.
- [11] A. Gil, J. Segura, J. A. G. Pertusa, and B. Soria, "Monte Carlo

simulation of 3-D buffered Ca²⁺ diffusion in neuroendocrine cells,” *Biophys. J.*, vol. 78, no. 1, pp. 13–33, 2000, doi: DOI:10.1016/s0006-3495(00)76569-6.

- [12] H. Casanova *et al.*, “The virtual instrument: Support for grid-enabled mcell simulations,” *Int. J. High Perform. Comput. Appl.*, vol. 18, no. 1, pp. 3–17, 2004, doi: DOI:10.1177/1094342004041290.
- [13] E. D. Andresta, N. A. Wibowo, and A. Sutresno, “Investigasi Pengaruh Jarak Celah Sinapsis dengan menggunakan Metode Monte Carlo,” *J. Fis. dan Apl.*, vol. 16, no. 3, p. 111, 2019, [Online]. Available: <https://iptek.its.ac.id/index.php/jfa/article/view/16%283%2901>
- [14] “DIFFUSION COEFFICIENTS OF MRI CONTRAST AGENTS IN CARTILAGE 46th Annual Meeting , Orthopaedic Research Society , March 12-15 , 2000 , Orlando , Florida,” p. 7021, 2000, [Online]. Available: <https://www.ors.org/transactions/46/0128.pdf>
- [15] F. R. Kuntari, S. Pranoto, K. A. Tiswati, and A. Sutresno, “Studi Proses Difusi melalui Membran dengan Pendekatan Kompartemen,” *J. Fis. dan Apl.*, vol. 15, no. 2, p. 62, 2019, [Online]. Available: <https://iptek.its.ac.id/index.php/jfa/article/view/4617>

